

Trigonometry Lesson #2: The Sine Law

Trigonometry in Acute Angled and Obtuse Angled Triangles

In the last lesson, we reviewed trigonometry in right triangles using SOHCAHTOA.

In the next three lessons, we focus on solving triangles which are not right angled and in which SOHCAHTOA is not valid.

In the next section of work we will determine the side of an acute angled triangle by

- i) splitting it in two right triangles and using SOHCAHTOA as in Class Ex. #1
- ii) using the Sine Law as in Class Ex. #2



Triangle ABC has three acute angles. Use SOHCAHTOA to determine the length of BC. Work to three decimal places and answer to two decimal places.

means less than 90°

SH $\sin 20 = \frac{z}{12.5}$
 $12.5 \sin 20 = z$
 $z = 4.275$

CA $\cos 20 = \frac{x}{12.5}$
 $12.5 \cos 20 = x$
 $x = 11.746$

TA $\tan 55 = \frac{y}{11.746}$
 $11.746 \tan 55 = y$
 $y = 16.775$

$BC = 4.275 + 16.775 = 21.05 \text{ cm}$


Copyright © by Absolute Value Publications. This book is NOT covered by the Cancopy agreement.

$a^2 + b^2 = c^2$
 SOHCAHTOA
 right angle triangles

A New Notation

Often, in trigonometry, it is convenient to use the following notation.

In triangle ABC , represent
 the length of the side opposite angle A by a ,
 the length of the side opposite angle B by b ,
 and the length of the side opposite angle C by c .




The Sine Law

In every triangle ABC , $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$ or $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$.

Proof of the Sine Law

The diagrams show the same triangle ABC placed with base AB on the x -axis.
 In diagram i) the origin is at A , and in diagram ii) the origin is at B .
 The line CD is drawn perpendicular to AB .



Complete the following work to show that $\frac{a}{\sin A} = \frac{b}{\sin B}$.

In i) $\sin A = \frac{CD}{AC} = \frac{CD}{b}$ In ii) $\sin B = \frac{CD}{BC} = \frac{CD}{a}$
 $CD = b \sin A$ $CD = a \sin B$

It follows that $b \sin A = a \sin B$

Dividing both sides by $\sin A \sin B$ gives the result

Repeating the work above with AC placed on the x -axis would give the result $\frac{a}{\sin A} = \frac{c}{\sin C}$.

Hence $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$ or $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$.

finding a side

finding an angle



To use the sine law, we need to know **three** pieces of information. This information must include both numerator and denominator of one of the three fractions, i.e. we need to know an angle and the measure of its opposite side.



Triangle ABC from Class Ex. #1 is shown. Use the sine law to calculate the length of BC, and compare your answer to the SOHCAHTOA method.

Finding a side

$$\frac{a}{\sin A} = \frac{c}{\sin C}$$

$$\frac{a}{\sin 75} = \frac{12.50}{\sin 35}$$

$$a = \frac{12.50 \sin 75}{\sin 35}$$

$$a = 21.05$$


Use the sine law in the triangle shown to determine the measure of $\angle ACB$ to the nearest degree.

Finding an angle

$$\frac{\sin A}{a} = \frac{\sin B}{b}$$

$$\frac{\sin 110}{9.4} = \frac{\sin B}{36.8}$$

$$0.5998 = \sin B$$

$$\sin^{-1}(0.5998) = B$$

$$B = 36.8$$

triangles add up to 180°
180 - 110 - 36.8 = 33°



A surveyor measures a base line PQ to be 440 m long. He takes measurements of a landmark R from P and Q, and finds that $\angle QPR = 46^\circ$ and $\angle PQR = 75^\circ$. Calculate the perimeter of $\triangle PQR$ to the nearest metre.

outsides add together

$$\frac{q}{\sin 75} = \frac{440}{\sin 59}$$

$$q = \frac{440 \sin 75}{\sin 59}$$

$$q = 495.8$$

$$\frac{p}{\sin 46} = \frac{440}{\sin 59}$$

$$p = \frac{440 \sin 46}{\sin 59}$$

$$p = 369.3$$

$$180 - 46 - 75 = 59$$

perimeter $q + p + r$
495.8 + 369.3 + 440 = 1305m




Complete Assignment Questions #1 - #9

Copyright © by Absolute Value Publications. This book is NOT covered by the Copyright agreement.




lac, 2ac, 3, 4 $\frac{1}{6}$

Assignment

1. Use the Sine Law to determine the length of the indicated side to the nearest tenth.




2. Use the Sine Law to determine the measure of the indicated angle to the nearest degree.




3. In $\triangle ABC$, angle $A = 49^\circ$, angle $B = 57^\circ$, and $a = 8$. Calculate b to the nearest tenth.

4. In $\triangle LMN$, angle $LMN = 114^\circ$, $LM = 123$ mm, and $MN = 88$ mm. Calculate $\angle LMN$, to the nearest degree.



5. P and Q are two bases for a mountain climb. PQ is 600 m and QR is a vertical stretch of a rock face. The angle of elevation of Q from P , (i.e. angle QPS) is 31° . The angle of elevation of R from P is 41° .




a) Mark these measurements on the diagram and state the measures of angle RPQ and angle PRQ .

b) Use the sine law in $\triangle PQR$ to calculate the height of the vertical climb, QR , to the nearest metre.

6. Consider the triangle shown.

- a) Use the sine law to calculate the lengths of the other two sides of the triangle to the nearest hundredth of a metre.



- b) Three students are trying to determine the area of the triangle in the diagram. Each student is given a different formula with which to determine the area. The area of the triangle is 53.3 m².

Show how each student arrived at this answer.

Student #1: Draw a vertical line to represent the height of the triangle and use the formula $A = \frac{1}{2}bh$, where b is the length of the base and h is the vertical height.

Student #2: Calculate the perimeter of the triangle and use Heron's formula $A = \sqrt{s(s-a)(s-b)(s-c)}$, where a , b , and c are the lengths of the three sides and s is the semi-perimeter of the triangle.


Student #3: Use the formula $A = \frac{1}{2}ab \sin C$, where a and b are the lengths of two sides and angle C is the contained angle between the sides a and b .

Multiple Choice

7. In triangle PQR , angle $P = 20^\circ$, angle $R = 150^\circ$, and $QR = 6$ m. The length of PQ is
- A. 4.1 m
 - B. 8.8 m
 - C. 15.2 m
 - D. 17.3 m
8. In $\triangle ABC$, $\angle A = 30^\circ$, $BC = 10$ units, and $AC = 15$ units. If $\angle B$ is acute-angled, then $\angle C$ is
- A. 19.4°
 - B. 48.6°
 - C. 101.4°
 - D. 130.6°

Numerical Response

9. From a point A , level with the foot of a hill, the angle of elevation of the top of the hill is 16° . From a point B , 950 metres nearer the foot of the hill, the angle of elevation of the top is 35° . The height of the hill, DC , to the nearest metre, is _____.



(Record your answer in the numerical response box from left to right.)

--	--	--	--

Answer Key

1. a) 12.4 cm b) 5.5 m c) 9.0 mm 2. a) 54° b) 44° c) 35°
 3. 8.9 4. 25°
 5. a) $10^\circ, 49^\circ$ b) 138 m 6. a) 9.52 m and 12.36 m
 7. B 8. C 9.

4	6	1	
---	---	---	--

Copyright © by Absolute Value Publications. This book is **NOT** covered by the Cancopy agreement.

Copyright © by Absolute Value Publications. This book is **NOT** covered by the Cancopy agreement.