

Quadratic Functions and Equations Lesson #1: Connecting Zeros, Roots, and x-intercepts

Function and Function Notation


Recall the following information from previous math courses:

Function

A functional relation, or **function**, is a special type of relation in which each element of the domain is related to exactly one element of the range. If any element of the domain is related to more than one element of the range, then the relation is not a function.

Function Notation

Under a function, f , the image of an element, x , in the domain is denoted by $f(x)$, which is read "f of x".



Consider a function f defined by the formula $f(x) = 4x - 8$.

The notation $f(x) = 4x - 8$ is called **function notation**.

We can show that, under the function f , the image of 5 is 12. We write $f(5) = 12$.

function notation

$$\begin{aligned} f(x) &= 4x - 8 \\ f(5) &= 4(5) - 8 \\ f(5) &= 12 \end{aligned}$$

equation of graph of function

$$\begin{aligned} y &= 4x - 8 \\ y &= 4(5) - 8 \\ y &= 12 \end{aligned}$$



We can also show that, under the function f , the image of 2 is 0. We write $f(2) = 0$.

function notation

$$\begin{aligned} f(x) &= 4x - 8 \\ f(2) &= 4(2) - 8 \\ f(2) &= 0 \end{aligned}$$

$x \quad y \quad (2, 0)$

equation of graph of function

$$\begin{aligned} y &= 4x - 8 \\ y &= 4(2) - 8 \\ y &= 0 \end{aligned}$$


We can say:

"The zero of the function $f(x) = 4x - 8$ is 2." "The root of the equation $y = 4x - 8$ is 2."

Zero(s) of a Function

A **zero of a function** is a value of the independent variable which makes the value of the function equal to zero. Zero(s) of a function can be found by solving the equation $f(x) = 0$.

x



Find the zero of the function f where $f(x) = 7x - 21$.

$$\begin{aligned} 0 &= 7x - 21 && f(x) = 0 \\ +21 & && \\ \hline 21 &= 7x && \\ \hline 3 &= x && \boxed{x=3} \end{aligned}$$

y
 x -int
zero
root

Copyright © by Absolute Value Publications. This book is **NOT** covered by the Cancopy agreement.

Investigation #1 *Connecting Roots, x-intercepts, and Zeros in a Linear Relation*

- a) The graph of $y = 2x - 6$ is shown. Determine the x -intercept of the graph algebraically and graphically.

$$\begin{aligned} \text{X-int, } y=0 & \quad 0 = 2x - 6 \\ & \quad \quad \quad x = 3 \end{aligned}$$

- b) Determine the root of the equation $2x - 6 = 0$.


$$\begin{aligned} 2x - 6 &= 0 \\ 2x &= 6 \\ x &= 3 \end{aligned}$$

- c) State the connection between the x -intercepts of the graph of $y = 2x - 6$ and the roots of the equation $2x - 6 = 0$.

Same

- d) Consider the function $f(x) = 2x - 6$. What is the zero of the function?

- e) What is the connection between the x -intercepts of the graph of $y = 2x - 6$, the roots of the equation $2x - 6 = 0$, and the zero of the function $f(x) = 2x - 6$?



Investigation #2 *Connecting Roots, x-intercepts, and Zeros in a Quadratic Relation*


- a) Jacques is determining the roots of the equation $x^2 - x - 6 = 0$.
 He wrote the equation in factored form and used the Zero Product Law to determine the roots of the equation.

ex. $(x-3)(x+2) = 0$
 $x = 3, -2$

Complete his work to solve for x .

$$\begin{array}{l} x^2 - x - 6 = 0 \\ (x-3)(x+2) = 0 \\ x = 3, -2 \end{array} \quad \begin{array}{l} x | + \\ -6 | -1 \end{array}$$

- b) The graph of $y = x^2 - x - 6$ is shown.
 State the x -intercepts of the graph and mark them on the grid.



- c) i) State the connection between the x -intercepts of the graph of $y = x^2 - x - 6$ and the roots of the equation $x^2 - x - 6 = 0$.

same

- ii) Explain the connection between the factors of $x^2 - x - 6$ and the roots of the equation $x^2 - x - 6 = 0$.

- d) Consider the function $g(x) = x^2 - x - 6$. Determine the zeros of the function.

c) Consider the function $g(x) = x^2 - x - 6$. Determine the zeros of the function.

e) State the connection between

- the **x-intercepts** of the graph of $y = x^2 - x - 6$
- the **roots** of the equation $x^2 - x - 6 = 0$, and
- the **zeros** of the function $g(x) = x^2 - x - 6$

all the same

Copyright © by Absolute Value Publications. This book is **NOT** covered by the Cancopy agreement.

270 Quadratic Functions and Equations Lesson #1: *Connecting Zeros, Roots, and x-intercepts*



a) Fill in the blanks in the following statement regarding the function with equation $y = f(x)$.

“The zeros of the function, the x-int of the graph of the function, and the roots of the corresponding equation $y = 0$, are the same numbers.”

b) The graph of $f(x) = x^2 - x - 6$ is shown. Fill in the blanks.

	The graph of	The function	The equation
	$f(x) = x^2 - x - 6$	$f(x) = x^2 - x - 6$ $= (x + 2)(x - 3)$	$x^2 - x - 6 = 0$
	has x-intercepts	has zeros	has roots
	$x = \underline{-2}$ and $x = \underline{3}$	$\underline{-2}$ and $\underline{3}$	$x = \underline{-2}$ and $x = \underline{3}$
	with y-intercept		
$y = \underline{-6}$			



Consider the equation $2x^2 - 7x + 3 = 0$.




a) Describe how the **zero feature** of a graphing calculator can be used to determine the roots of the equation.

b) Use a graphing calculator to determine the roots of the



b) Use a graphing calculator to determine the roots of the equation and sketch the graph on the grid provided.

c) Use the x -intercepts of the graph of $y = 2x^2 - 7x + 3$ to factor the expression $2x^2 - 7x + 3$.



d) What are the zeros of the function $f(x) = 2x^2 - 7x + 3$?

Copyright © by Absolute Value Publications. This book is **NOT** covered by the Cancopy agreement.



Finding Zeros of a Function

To find the zeros of a function, $f(x)$, either

- substitute zero for $f(x)$ and find the roots of the resulting equation
- or
- graph the function and determine the x -intercepts of the graph

Finding the Roots of an Equation by Factoring

Finding the roots of a single variable equation may involve factoring. Except in the case of a linear equation, set the equation to zero before factoring.

Recall the following techniques for factoring common factors, difference of two squares, trinomials of the form $x^2 + bx + c = 0$, and trinomials of the form $ax^2 + bx + c = 0$.



Find the roots of the following equations.

a) $x^2 + 8x = 33$

$x^2 + 8x - 33 = 0$
 $(x+11)(x-3) = 0$
 $x = -11, 3$
 roots are -11 and 3

b) $6(4x + 5)(x - 3) = 0$

$4x + 5 = 0$ $x - 3 = 0$
 $4x = -5$ $x = 3$
 $x = -\frac{5}{4}$
 roots are $-\frac{5}{4}$ and 3

c) $2x^2 - 8 = 0$

$2(x^2 - 4) = 0$
 $2(x-2)(x+2) = 0$
 $x = -2, 2$
 ± 2
 roots are -2 and 2



For the following functions



For the following functions

- i) find the zeros ii) find the y-intercept of the graph of the function

a) $f(x) = 5x^2 + 15x - 20$

i) $5(x^2 + 3x - 4) = 0$

$5(x+4)(x-1) = 0$

$x = -4, 1$

zeros are -4 and 1

ii) y-int when $x=0$

$f(0) = 5(0)^2 + 15(0) - 20$

$f(0) = -20$

y-int = -20

b) $f(x) = 3x^2 - 11x + 10$

$3x^2 - 6x - 5x + 10 = 0$

$3x(x-2) - 5(x-2) = 0$

$(x-2)(3x-5) = 0$

$x = 2, \frac{5}{3}$

zeros are 2 and $\frac{5}{3}$

ii) y-int, $x=0$

$f(0) = 3(0)^2 - 11(0) + 10$
 $= 10$

y-int = 10

c) $g(x) = 2x(2x+1)$

$2x(2x+1) = 0$

$x = 0, -\frac{1}{2}$

zeros are 0 and $-\frac{1}{2}$

ii) y-int, $x=0$

$g(0) = 2(0)(2(0)+1) = 0$

$y = 0$

y-int = 0

Copyright © by Absolute Value Publications. This book is NOT covered by the Cancopy agreement.


272 Quadratic Functions and Equations Lesson #1: Connecting Zeros, Roots, and x-intercepts



For each of the following functions, use a graphing calculator to

- sketch the graph of the function
- find the zeros of the function as exact values
- write the function in factored form

a) $f(x) = 3x^2 + 4x - 7$




zeros: $-\frac{7}{3}, 1$

$= 3(x + \frac{7}{3})(x - 1)$

$= (3x - 7)(x - 1)$

b) $g(x) = 4x^3 - 7x^2 - 4x + 7$




zeros: $-1, 1, \frac{7}{4}$

$= 4(x+1)(x-1)(x-\frac{7}{4})$

$= (x+1)(x-1)(4x-7)$


Complete Assignment Questions #1 - #11

Assignment



Assignment

1. The graph of a function, f , is shown. The x and y -intercepts of the graph are integers.
- State the x and y -intercepts of the graph.
 - State the zeros of the function f .



2. Find the roots of the following equations.
- $2x(x + 3) = 0$
 - $2x^2 - 10x + 12 = 0$
 - $x^3 + 8x^2 = 20x$
 - $4x^2 + 4x - 3 = 0$

Copyright © by Absolute Value Publications. This book is **NOT** covered by the Cancopy agreement.

Quadratic Functions and Equations Lesson #1: *Connecting Zeros, Roots, and x -intercepts* **273**

3. Find the zeros of the following functions.

a) $f(x) = \frac{x}{3} + 5$

b) $g(x) = 25x^2 - 64$

c) $P(x) = 3(2x - 5)(x + 1)$

d) $P(x) = x(x - 3)(2x + 1)$

4. In each of the following
- determine the zeros of the function
 - determine the y -intercept of the graph of the function

a) $f(x) = 5x^2 - 35x$

b) $f(x) = 3x(x^2 - 49)$

c) $f(x) = 2x^2 - x - 15$

d) $P(x) = 8x^2 + 14x - 15$

5. Use a graphing calculator to find the zeros (as exact values) of the following functions.

a) $f(x) = 18x^2 - 5x - 7$

b) $g(x) = 3x^3 - 11x^2 + 6x$




Copyright © by Absolute Value Publications. This book is **NOT** covered by the Cancopy agreement.


274 Quadratic Functions and Equations Lesson #1: *Connecting Zeros, Roots, and x-intercepts*

6. In each case, the graph of a function with $y = f(x)$ is shown. The x and y -intercepts of the graph are integers. Determine

- the zeros of the function
- the y -intercept of the graph of the function
- the equation of the function in factored form Bonus

1, 2, 3, 4, 6





7. Use a graphing calculator to write the equation in factored form.

a) $y = 2x^2 - 3x - 9$

b) $y = 5x^3 - 7x^2 - 21x - 9$

Multiple Choice 8. The zeros of the function $f(x) = 2(x - 3)(4x + 7)$ are

A. $3, -\frac{7}{4}$

B. $-3, \frac{7}{4}$

C. $0, 3, -\frac{7}{4}$

D. $2, 3, -\frac{7}{4}$

Copyright © by Absolute Value Publications. This book is **NOT** covered by the Cancopy agreement.

9. The roots of the equation $3x(x + 1) = 6$ are

A. $0, -1$

B. $2, 5$

C. $2, -1$

D. $-2, 1$

10. The least possible zero of the function $f(x) = 2x^3 - 7x^2 + 3x$ is

A. 0

B. $\frac{1}{2}$

- A. 0
- B. $\frac{1}{2}$
- C. 3
- D. -3

Numerical Response 11. The y-intercept of the graph of the function $f(x) = (x + 4)(3 - 2x)(x + 1)$, to the nearest whole number, is _____ .

(Record your answer in the numerical response box from left to right.)

--	--	--	--

Answer Key

1. a) x-intercepts are -6, 4 and y-intercept is -24. b) -6, 4

2. a) -3, 0 b) 2, 3 c) -10, 0, 2 d) $-\frac{3}{2}, \frac{1}{2}$

3. a) -15 b) $-\frac{8}{5}, \frac{8}{5}$ c) $-1, \frac{5}{2}$ d) $-\frac{1}{2}, 0, 3$

4. a) i) 0, 7 ii) 0 b) i) -7, 0, 7 ii) 0
 c) i) $-\frac{5}{2}, 3$ ii) -15 d) i) $-\frac{5}{2}, \frac{3}{4}$ ii) -15

5. a) $-\frac{1}{2}, \frac{7}{9}$ b) $0, \frac{2}{3}, 3$

6. a) zero: 3 b) zeros: -2, 4 c) zeros: -2, 3, 4
 y-intercept: 6 y-intercept: -16 y-intercept: 24
 $f(x) = -2(x - 3)$ $f(x) = 2(x + 2)(x - 4)$ $f(x) = (x + 2)(x - 3)(x - 4)$

7. a) $y = (2x + 3)(x - 3)$ b) $y = (5x + 3)(x - 3)(x + 1)$

8. A 9. D 10. A 11.

1	2		
---	---	--	--

Copyright © by Absolute Value Publications. This book is **NOT** covered by the Cancopy agreement.

*Copyright © by Absolute Value Publications. This book is **NOT** covered by the Cancopy agreement.*